

CONFIDENTIAL		SEE BOTTOM OF PAGE FOR SPECIAL CONTROLS, IF ANY	
CONFIDENTIAL INFORMATION REPORT		This material contains information affecting the National Defense of the United States within the meaning of the Espionage Laws, Title 18, U.S.C. Secs. 793 and 794, the transmission or revelation of which in any manner to an unauthorized person is prohibited by law.	
PREPARED AND DISSEMINATED BY CENTRAL INTELLIGENCE AGENCY			
COUNTRY USSR			
SUBJECT [] Medical Installation [] []	DATE DISTRIBUTED 26 Dec 1956		
	NO. OF PAGES 1	NO. OF ENCLS.	
	SUPPLEMENT TO REPORT #		
THIS IS UNEVALUATED INFORMATION			

25X1

25X1

25X1

25X1

25X1

- end -

~~CONFIDENTIAL~~

DISTRIBUTION	STATE	ARMY	NAVY	AIR															
--------------	-------	------	------	-----	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

NOFORN**NO DISSEM ABROAD****LIMITED**

LIMITED: Dissemination limited to full-time employees of CIA, AEC and FBI; and, within State and Defense, to the intelligence components, other offices requiring NIS elements, and other agencies with their immediate supporting staffs. Not to be disseminated to consultants, external agencies or persons.

Approved For Release 2008/01/25 : CIA-RDP80T00246A000500220001-1

Page Denied

Next 11 Page(s) In Document Denied

HOSPITAL

[] the wards in the Clinic Uglov, where the patients looked clean, content and well cared for []

[] Women as surgeons seem to dominate the surgical scene. []

one plump motherly soul of probably 57 proceeding with a gastric resection under local 25X1
anesthesia. The patient was a serene, alert woman of perhaps 60. One (female) assist-
ant physician [] constantly and rapidly injected around the edges of the wound 25X1
0.25 per cent procaine in 10 cc. syringefuls, keeping out of the way of the surgeon as
well as possible. Once the initial anesthetization had been carried out both surgery 25X1
and further anesthesia progressed simultaneously. About 2 liters of this "local"
solution are used for a gastric resection and 3 liters may be used for a pulmonary 25X1
resection! Doubtless a good deal of the procaine solution leaks out from the wound.
Epinephrine is added, 2 drops (1:1000) epinephrine to each 100 cc. The premedication 25X1
employed is 1 cc. 1 per cent morphine subcutaneously (10 mg.)

25X1

FOR OFFICIAL USE ONLY

Uglov prefers ether for open chest surgery but can do it under local with (preferably) oxygen under pressure.

[] operation in the clinic under endotracheal ether-oxygen.

25X1

Uglov sometimes uses 200 cc. blood with a gastric resection. He often uses placenta blood, but does not yet have cadaver blood. []

25X1

Grouping but not cross matching is carried out. Patients routinely get an electrolyte solution, called "antishock," according to the following formula:

Sodium chloride	3.75
Glucose	25.
Morphine	0.0075
Magnesium	1.5
"Revanole"(antiseptic)	0.005
Distilled water	500.

[] this can be stored and used over a period of many months.

25X1

Uglov combines hypothermia with "artificial hibernation" sometimes.

#

25X1

[] of the 3500 medical students in the Leningrad school 70 per

25X1

cent are women. []

25X1

25X1

FOR OFFICIAL USE ONLY

Page Denied

Next 14 Page(s) In Document Denied